

# **Spectral-Fluorescence Properties of the $\text{Sr}_3\text{LaMe}_3\text{O}_{12}$ ( Me = Nb, Ta ) Crystals Activated by $\text{Nd}^{3+}$ Ions**

M. Kh. Ashurov and O.K. Alimov  
“Phonon” Scientific Industrial Association  
Yusupov str. 50  
700054 Tashkent, Uzbekistan

A.Sh. Yarmuhamedov and H.T. Shodiev  
Samarkand State University  
University blvd. 15  
703004 Samarkand

In this report, the results on spectral and kinetic investigation obtained by site- selective laser spectroscopy of the  $\text{Sr}_3\text{LaNb}_3\text{O}_{12}$  (SLN) and the  $\text{Sr}_3\text{LaTa}_3\text{O}_{12}$  (SLT) crystals doped with  $\text{Nd}^{3+}$  ions are presented. The  $\text{Nd}^{3+}$  ions concentrations in studied crystals were 0,5 – 5 weight %. Time resolved fluorescence spectra (  $^4\text{F}_{3/2} \rightarrow ^4\text{I}_{9/2}; ^4\text{I}_{11/2}$  ) and laser excitation spectra (  $^4\text{I}_{9/2} \rightarrow ^4\text{G}_{5/2}; ^2\text{G}_{7/2}$  ) were registered at 77 and 300K.

Results show the creation of the three types of centers of  $\text{Nd}^{3+}$ , A,B,C ,in the SLN crystal and only two types of centers in SLT, B' and C', similar to B and C centers in SLN.

Neodimium ion was found to occupy mainly two positions with different surroundings which correspond to the B and C centers. Also, the crystal field fluctuations result in the inhomogeneous distribution of energies of the  $[^4\text{F}_{3/2}(1) \rightarrow ^4\text{I}_{9/2}(1)]$  electronic transition of the  $\text{Nd}^{3+}$  ions with spectral broadening equal to  $\Delta\nu_{\text{inhom}} = 17\text{cm}^{-1}$  for both B, B' and C, C' centers. The magnitude of homogeneous broadening  $[\delta_{\text{hom}}(\nu) = 27\text{cm}^{-1}]$  of the  $^4\text{F}_{3/2}(1) \rightarrow ^4\text{I}_{9/2}(5)$  transition of A-center is determined at 300 K. There are proposed different models of the  $\text{Nd}^{3+}$  optical centers formation in used crystals matrices. It is shown that B, B' and C, C' optical centers have  $\text{C}_{3v}$  point group symmetry. Energy level diagrams of Stark-splitting of spectral terms  $^4\text{I}_{9/2}; ^4\text{I}_{11/2}; ^4\text{F}_{3/2}; ^4\text{G}_{5/2}; ^2\text{G}_{7/2}$  of the A, B, B', and C, C' centers are obtained. By means of fluorescence line narrowing the kinetics of fluorescence decay and lifetimes of excited  $^4\text{F}_{3/2}$  state for the A, B, B', and C, C' - centers are measured. The  $\text{Nd}^{3+}$  -  $\text{Nd}^{3+}$  inter-center energy transfer in SLN crystal was observed.